


Vitamin D3 1,000 IU



CLINICAL APPLICATIONS

- Maintains Bone and Dental Health
- Supports Calcium Absorption and Balance
- Boosts Immune Activity
- Maintains Normal Cardiometabolic Function, and Supports Blood Sugar Balance Already Within Normal Levels
- Helps Support Musculoskeletal Strength and Comfort

ESSENTIAL VITAMINS

What Is Vitamin D?

Vitamin D is a steroid vitamin, a group of fat-soluble pro-hormones that are best known for the role they play in supporting bone health and aiding in the absorption of calcium and phosphate from the gastrointestinal tract. A growing body of research highlights its important role in supporting other body systems, including cardiovascular and blood sugar balance, musculoskeletal strength, and neurological and immune function, enabled by its ability to target over 200 different genes throughout the body. At the same time, deficiency and insufficiency of this important nutrient has reached epidemic proportions around the world, making the achievement of optimal levels more important than ever to overall health and wellbeing.

Overview

Known as the sunshine vitamin, one of the key roles of vitamin D is maintaining serum calcium and phosphorous balance. Our bodies make vitamin D by converting vitamin D2 to D3, or cholecalciferol (the active form), when exposed to sunlight. D3 is also the form the body derives from dietary cholesterol. When calcium and phosphorus levels dip in the body, parathyroid hormone (PTH) is released to increase vitamin D conversion to the active form. D3 is then metabolized to calcitriol, a steroid hormone that helps regulate a variety of genes through the vitamin D receptor (VDR). While vitamin D is available in both forms as supplements, studies have found vitamin D3 is the preferable form, as it has been found to maintain active vitamin D levels for a longer period of time. This vitamin D formulation is delivered as D3 (cholecalciferol) for optimal use by the body and is available in 1,000 and 5,000 IU capsules to meet a variety of individual needs.

Vitamin D Depletion[†]

While it has long been assumed that the majority of the population achieves adequate levels of vitamin D through exposure to the sun, the biosynthesis of the nutrient is affected by time of day, seasons, location, smog/pollution, clothing and sunscreen. In addition, those with darker skin require more exposure to the sun to attain adequate levels. These factors all contribute to the insufficiency seen in a growing portion of the population. It takes about 48 hours for vitamin D to be absorbed from the skin into the body and washing skin during that time period can interfere with absorption into the blood. Inadequate intake or levels of cholesterol can also inhibit the adequate production of the nutrient. Depleted levels of vitamin D may interfere with the normal development of teeth and bones, normal cell growth, and contribute to poor regulation of the immune and nervous systems. Certain medications have been found to contribute to deficiency of vitamin D, including some anti-seizure medications, bile acid sequestrants, oral corticosteroids, and weight loss medications, which bind fats.¹

Bone and Dental Health[†]

Numerous studies have highlighted the importance of vitamin D to maintaining healthy bone density. In one 2013 study, 52 overweight men and women with suboptimal vitamin D levels were given either 7,000 IU of cholecalciferol daily or a placebo for 26 weeks. The vitamin D group significantly increased vitamin D levels in the blood and improved biomarkers of bone health.^{2,3}

[†] These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Cardiometabolic Health and Blood Sugar Balance[†]

In light of the role that vitamin D plays in a variety of tissues and body systems, it comes as no surprise that much research from recent years has underscored its contribution to cardiometabolic wellness.⁴ There is accumulating evidence that calcitriol helps strengthen cardiomyocyte function, vascular smooth muscle cells, and the vascular endothelium. Low levels of 25-hydroxyvitamin D are associated with diminished cardiovascular health.⁵ In a recent study of 222 participants, slow coronary function was significantly higher in those with insufficient blood levels of the vitamin.⁶ Optimal vitamin D status has also been linked with a 40% cardiovascular system protective effect⁷ as well as maintenance of normal blood pressure levels.⁸ Vitamin D has also been shown to support normal blood sugar metabolism.^{9,10}

Immune Health and Modulation[†]

One of the more profound functions of vitamin D is its ability to modulate immunity. An important recent study suggested that improving vitamin D status significantly affects the expression of genetic pathways linked to immune activity.¹¹ Vitamin D has been shown to boost the immune response by up-regulating specific genes that increase cellular production of natural compounds that protect us against pathogens.¹²

Musculoskeletal Comfort[†]

Numerous studies also point to the key role of vitamin D in supporting musculoskeletal strength and comfort.¹³ In one study among 62 adult patients with nonspecific musculoskeletal discomfort, over 95% had vitamin D deficiency and responded to replenishment of vitamin D. Moderate deficiency of vitamin D has also been shown to predict knee discomfort over a five-year period and hip discomfort over two years.¹⁴

Directions

1 capsule per day or as recommended by your health care professional.

Does Not Contain

Gluten, yeast, artificial colors or flavors.

Cautions

If you are pregnant or nursing, consult your physician before taking this product.

Supplement Facts^{V4}

Serving Size 1 Capsule
Servings Per Container 180

	Amount Per Serving	% Daily Value
Vitamin D (D3 as Cholecalciferol)	25 mcg (1,000 IU)	125%

Other Ingredients: Microcrystalline Cellulose, Hypromellose (Natural Vegetable Capsule), Magnesium Stearate, Silicon Dioxide and Stearic Acid.

ID# 108180 180 Capsules

[†] These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

References

1. <http://umm.edu/health/medical/altmed/supplement/vitamin-d>
2. Wamberg L, Pedersen SB, Richelsen B, Rejnmark L. The effect of high-dose vitamin d supplementation on calciotropic hormones and bone mineral density in obese subjects with low levels of circulating 25-hydroxyvitamin d: results from a randomized controlled study. *Calcif Tissue Int*. 2013 Jul;93(1):69-77.
3. Cauley JA, Lacroix AZ, Wu L, Horwitz M, Danielson ME, Bauer DC, Lee JS, Jackson RD, Robbins JA, Wu C, Stanczyk FZ, LeBoff MS, Wactawski-Wende J, Sarto G, Ockene J, Cummings SR. Serum 25-hydroxyvitamin D concentrations and risk for hip fractures. *Ann Intern Med*. 2008 Aug 19;149(4):242-50.
4. Wang C. Role of vitamin d in cardiometabolic diseases. *J Diabetes Res*. 2013;2013:243934. doi:10.1155/2013/243934
5. Zittermann A, Koerfer R. Vitamin D in the prevention and treatment of coronary heart disease. *Curr Opin Clin Nutr Metab Care*. 2008 Nov;11(6):752-7.
6. Oz F, Cizgici AY, Oflaz H, Elitok A, Karaayvaz EB, Mercanoglu F, Bugra Z, Omer B, Adalet K, Oncul A. Impact of vitamin D insufficiency on the epicardial coronary flow velocity and endothelial function. *Coron Artery Dis*. 2013 May 20.
7. Ng LL, Sandhu JK, Squire IB, Davies JE, Jones DJ. Vitamin D and prognosis in acute myocardial infarction. *Int J Cardiol*. 2013 Feb 13.
8. Martini, L.A. and Wood, R.J. Vitamin D and blood pressure connection: update on epidemiologic, clinical, and mechanistic evidence. *Nutr Rev*. 2008; 66(5):291-297.
9. Teegarden D, Donkin SS. Vitamin D: emerging new roles in insulin sensitivity. *Nutr Res Rev*. 2009 Jun;22(1):82-92.
10. Chiu KC, Chu A, Go VL, Saad MF. Hypovitaminosis D is associated with insulin resistance and beta cell dysfunction. *Am J Clin Nutr*. 2004 May;79(5):820-5.
11. Hossein-nezhad A, Spira A, Holick MF. Influence of vitamin D status and vitamin D3 supplementation on genome wide expression of white blood cells: a randomized double-blind clinical trial. *PLoS One*. 2013;8(3):e58725.
12. Cannell JJ, Vieth R, Umhau JC, Holick MF, Grant WB, Madronich S, Garland CF, Giovannucci E. Epidemic influenza and vitamin D. *Epidemiol Infect*. 2006 Dec;134(6):1129-40.
13. Abbasi M, Hashemipour S, Hajmanuchehri F, Kazemifar AM. Is vitamin D deficiency associated with non-specific musculoskeletal pain? *Glob J Health Sci*. 2012 Nov 11;5(1):107-11.
14. Laslett LL, Quinn S, Burgess JR, Parameswaran V, Winzenberg TM, Jones G, Ding C. Moderate vitamin D deficiency is associated with changes in knee and hip pain in older adults: a 5-year longitudinal study. *Ann Rheum Dis*. 2013 Apr 17.